

Solvent effect on the volume of activation and volume of the Diels-Alder reaction

Kiselev V., Konovalov A., Asano T., Kashaeva E., Iskhakova G., Shihab M., Medvedeva M.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

Volumes of activation, ΔV^\ddagger , and reaction, ΔV , partial molar volumes, V , and enthalpies of solution, $\Delta_{\text{sol}}H$, were determined for tetracyanoethylene, cyclopentadiene, 1,3-butadiene, trans, trans-1,4-diphenyl-1,3-butadiene and their Diels-Alder adducts at 25°C in some solvents of π - and n-donor type. The values of the activation and reaction volumes were exceptionally small in the former type of solvents. Large solvent effects on $\Delta_{\text{sol}}H_{\text{TCNE}}$ (up to 26 kJ mol⁻¹), V_{TCNE} , ΔV and δV^\ddagger (up to 11 cm³mol⁻¹) were observed in aromatic solvents and these values are linearly correlated with each other. Poor correlations were found for n-donor solvents, but a linear dependence between ΔV and $\delta_r nH$ (enthalpy of reaction) was obtained for all solvents. © 2001 John Wiley & Sons, Ltd.

<http://dx.doi.org/10.1002/poc.398>

Keywords

Solvent effect, Tetracyanoethylene, Volume of activation, Volume of reaction